arXiv:1303.4864v1 [quant-ph] 20 Mar 2013

Coherent decay of dressed states in Jaynes-Cummings model with a common bath

Zhihai Wang and D. L. Zhou*
Beijing National Laboratory for condensed matter physics,
Institute of Physics, Chinese Academy of Sciences, Beijing, 100190, China

We propose an experimental scheme to realize the Jaynes-Cummings model where the two-level atom and the cavity photon share a common bath. We find that the cavity photon will have a dramatically slow decay rate, and the vacuum Rabi splitting in the transmission spectrum shows an obvious asymmetric character. These two results can be explained based on the different decay rates of two dressed states. Quantum coherence enhances the decay rate of one dressed state but suppresses the decay rate of the other.

Introduction.— In nature, any quantum system inevitably interacts with its environment [1]. In addition, a quantum measurement on a quantum system must introduce some measurement device [2]. Thus, to understand the fundamental concepts in quantum mechanics, we have to study quantum open systems. Furthermore, to control and manipulate an open quantum system is a central task in quantum information [3–5].

In literature, extensive investigations on quantum open system have been done. In particular, a spin-boson model is used as a typical paradigm in studying quantum open system [6, 7]. In the spin-boson model, a single two-level spin is regarded as the system, and the environment is composed by many different bosonic modes. Here, we will investigate the atom-cavity coupled system with a common bath. We describe the system by the Jaynes-Cummings model(JCM) [8], and aim to show how a common bath may essentially change the properties of the system, particularly on the effects induced by quantum coherence.

A central task in experiment is how to realize a common bath shared by the atom and the cavity. Motivated by the recent experimental setting in Ref. [9], we propose a scheme as shown in Fig. 1, in which a two-level atom is located in a dissipative cavity, and the leaky light from the cavity is reflected back to interact with the atom by four mirrors. In our scheme, the environment is composed by the light modes in the ring cavity formed by the four mirrors, which is shared by the two-level atom and the cavity mode. When the atom is nearly resonant with the cavity mode, the photons emitted by the atom may be absorbed by the cavity mode assisted by the environment, and vice versa. These feedback processes assisted by the common bath will play an important role in our scheme.

Theoretically, the master equation is widely used to investigate the non-equilibrium dynamics of the open system [10–16]. However, most of the previous investigations in the cavity-QED system regard the cavity dissipation and the spontaneous emission of the atom as two independent processes. However, such treatment is no longer valid in our scheme because of the feedback processes discussed above. Hence we take into account the coherence effect of the two-level atom and the cavity mode by reformulating the traditional master equation under the secular approximation [17], which is confirmed by the numerical results.



FIG. 1. (Color online) The proposed experimental scheme. A two-level atom is located in a leaky cavity which supports a single mode electromagnetic field. The photons outside the cavity can be reflected by the mirrors and serve as a common bath shared by the atom and the cavity. The photon detector(PD) is applied to detect the photons in the cavity.

In this letter, we focus on the effect of the coherence between the atom and the cavity on the dynamical evolution and the transmission spectrum of the coupled system. We find that both of the atom and the cavity will finally decay to their ground states after several cycles of Rabi-like oscillations in the absence of the driven field. The atom-cavity coherence dramatically slows down the decay of the system. Furthermore, in the presence of a classical driven field, the vacuum Rabi splitting [18–25] in the transmission spectrum shows an obvious asymmetric character. These two exotic phenomena originate from the fact that the quantum coherence enhances the decay of symmetric dressed state while suppresses the antisymmetric dressed state.

Model.— We propose an experimental scheme as shown in Fig. 1. A two-level atom is located in a high finesse cavity which supports a single mode electromagnetic field. The light modes in the ring cavity formed by four mirrors construct the common bath shared by the cavity mode and the two-level atom.

The Hamiltonian of the global system can be written as the sum of three terms: $H = H_{JC} + H_B + H_I$, where

$$H_{JC} = \omega_c a^\dagger a + \frac{\omega_0}{2}\sigma_z + g(a^\dagger\sigma^- + a\sigma^+), \tag{1a}$$

$$H_B = \sum_i \omega_i b_i^\dagger b_i, \tag{1b}$$

$$H_I = \sum_i \kappa_i(ab_i^\dagger + b_i a^\dagger) + \sum_i \xi_i(\sigma^- b_i^\dagger + b_i\sigma^+). \tag{1c}$$

The first term H_{JC} is the Hamiltonian of our system, the JCM, which describes a two-level atom interacting with the single mode cavity photon under the rotating wave approximation. In Eq. (1a), ω_c is the frequency of the cavity mode, a is the annihilation operator of the mode. The two energy levels of the atom are denoted as $|g\rangle$ and $|e\rangle$, and ω_0 is the energy difference. The Pauli operators are defined as $\sigma_z \equiv |e\rangle\langle e| - |g\rangle\langle g|$, $\sigma^- \equiv |g\rangle\langle e|$, and $\sigma^+ \equiv |e\rangle\langle g|$.

The second term H_B describes the free terms of the photons in the ring cavity, which act as a bath in our scheme. In Eq. (1b), ω_i is the frequency of the i-th mode in the ring cavity and b_i is its annihilation operator.

The third term H_I describes the interactions between the system and the bath. In Eq. (1c), κ_i (ξ_i) is the coupling strength between the atom (the cavity photon) and the i-th mode of the bath.

Decay of a cavity photon. — To investigate the effect induced by the common bath, we study how a single cavity photon decays into the bath. In other words, when the initial state of the global system is $|\psi(0)\rangle = |1; g\rangle \otimes_i |0_i\rangle$, what is the average cavity photon number during the evolution of the system?

Notice that there are two decay channels for the cavity photon. First, the cavity photon can directly decay into the bath. Second, the cavity photon can be absorbed by the atom, then the atom emits a photon into the bath. An intuitive idea is to sum the effects of the two channels [19]:

$$\dot\rho = -i[H, \rho] + J_1(\omega_c)\mathcal{L}[a] + J_2(\omega_0)\mathcal{L}[\sigma^-], \tag{2}$$

where $\mathcal{L}[Q] = (2Q\rho Q^\dagger - Q^\dagger Q\rho - \rho Q^\dagger Q)$, and the spectrum functions $J_1(\omega)$ and $J_2(\omega)$ are defined as [6, 26]

$$J_1(\omega) = \pi\sum_j \kappa_j^2\delta(\omega - \omega_j), \tag{3a}$$

$$J_2(\omega) = \pi\sum_j \xi_j^2\delta(\omega - \omega_j). \tag{3b}$$

However, there are two important factors that are not taken into account in the above intuitive approach. Firstly, the interaction between the atom and the cavity photon leads to the dressed state of the atom and the cavity. The energy differences between two adjacent dressed states depend on the system parameters ω_c, ω_0 and g. Because, in the eye of the bath, the atom and the cavity photon are dressed, it is not valid to consider the decay channel in the bare states. Secondly, because the cavity mode and the atom share a common bath, the bath photon emitted by the atom can be reabsorbed by the cavity, and vice versa. The feedback processes are completely neglected in the intuitive method.

To take into account the above two factors, we adopt the picture of dressed states in the JCM. The ground state of H_{JC} is a product state $|G\rangle = |0; g\rangle$ with eigen-energy $E_0 = -\omega_c/2$. In the resonance case ($\omega_c = \omega_0$), the energies for the excited states are $E_n^\pm = (n - 1/2)\omega_c \pm g\sqrt{n}$, the corresponding eigen-vectors are the dressed states $|n, \pm\rangle = (\pm|n; g\rangle + |n-1; e\rangle)/\sqrt{2}$, which are coherent superpositions of the product states $|n; g\rangle$ and $|n-1; e\rangle$.

Applying the secular approximation [17], we re-derive the master equation in the dressed state picture, whose detailed calculation is given in the supplemental material [27]. The master equation we obtain is

$$\dot\rho_{cd} = -i(E_c - E_d)\rho_{cd} + \sum_{k,l}\gamma^{cdkl}\rho_{kl} \tag{4}$$

where $|c\rangle, |d\rangle, |k\rangle, |l\rangle$ are the dressed states of H_{JC} with the eigen-energies E_c, E_d, E_k and E_l respectively, $\rho_{cd} = \langle c|\rho|d\rangle$ and $\rho_{kl} = \langle k|\rho|l\rangle$ are the elements of the reduced density matrix for the atom-cavity system. In Eq. (4), $\gamma^{cdkl} \equiv \sum_{i=1}^4 \gamma_i$ with

$$\gamma_1 = -\sum_n\left[J_1(\omega_{kn})\delta_{dl}a^\dagger_{cn}a_{nk} + J_2(\omega_{kn})\delta_{dl}\sigma^+_{cn}\sigma^-_{nk} + \sqrt{J_1(\omega_{kn})J_2(\omega_{kn})}\delta_{dl}\left(a^\dagger_{cn}\sigma^-_{nk} + \sigma^+_{cn}a_{nk}\right)\right] \tag{5a}$$

$$\gamma_2 = J_1(\omega_{kc})a_{ck}a^\dagger_{ld} + J_2(\omega_{kc})\sigma^-_{ck}\sigma^+_{ld} + \sqrt{J_1(\omega_{kc})J_2(\omega_{kc})}\left(a_{ck}\sigma^+_{ld} + \sigma^-_{ck}a^\dagger_{ld}\right) \tag{5b}$$

$$\gamma_3 = -\sum_n\left[J_1(\omega_{ln})\delta_{ck}a^\dagger_{ln}a_{nd} + J_2(\omega_{ln})\delta_{ck}\sigma^+_{ln}\sigma^-_{nd} + \sqrt{J_1(\omega_{ln})J_2(\omega_{ln})}\delta_{ck}\left(a^\dagger_{ln}\sigma^-_{nd} + \sigma^+_{ln}a_{nd}\right)\right] \tag{5c}$$

$$\gamma_4 = J_1(\omega_{ld})a_{ck}a^\dagger_{ld} + J_2(\omega_{ld})\sigma^-_{ck}\sigma^+_{ld} + \sqrt{J_1(\omega_{ld})J_2(\omega_{ld})}\left(a_{ck}\sigma^+_{ld} + \sigma^-_{ck}a^\dagger_{ld}\right) \tag{5d}$$

In Eqs. (5), $\omega_{ij} = E_i - E_j$ is the energy difference between level i and j, and $A_{\alpha\beta} \equiv \langle\alpha|A|\beta\rangle$ is the matrix element of operator A in the dressed state representation of the JCM. In the above equations, the terms proportional to $\sqrt{J_1(\cdot)J_2(\cdot)}$ represent the contribution from the atom-cavity coherence.

Under the Ohmic dissipation, the spectrum functions $J_1(\omega)$ and $J_2(\omega)$ of the bath are expressed as

$$J_1(\omega) = 2\pi\alpha_1\omega\exp(-\omega/\omega_{c1}), \tag{6a}$$

$$J_2(\omega) = 2\pi\alpha_2\omega\exp(-\omega/\omega_{c2}), \tag{6b}$$

where $\alpha_{1(2)}$ are the dissipation coefficients and $\omega_{c1(c2)}$ denote the cutoff frequencies. Solving the master equation Eq. (4) numerically, we plot the curve of the average photons in the cavity $\langle a^\dagger a\rangle$ as a function of the evolution time in Fig. 2.



FIG. 2. (Color online) The average photon number in the cavity as a function of the evolution time t. The parameters are set as $\omega_c = \omega = 1$, $g = 0.1$, $\omega_{c1} = 5$, $\omega_{c2} = 8$, $\alpha_1 = 0.002$, $\alpha_2 = 0.001$. The cut off photon number in the numerical calculation is $N = 2$. The initial state is $|\psi(0)\rangle = |1; g\rangle \otimes_i |0_i\rangle$.

In Fig. 2, we also plot the results given by neglecting the contribution from the atom-cavity coherence, the results given by Eq. (2) as well as the numerical results which is obtained from the solution of the Schrödinger equation. It is shown that, the results given by our approach agree with the numerical results very well. Since we consider the case of Ohmic dissipation, J_1 and J_2 are not very sensitive to the change of the variable ω when $\omega \ll \omega_{c1(c2)}$. Therefore, the results given by neglecting the effect of the coherence nearly coincide with the results from Eq. (2).

It is shown in Fig. 2, as expected in the region of strong coupling, the Rabi-like oscillation occurs in the short time as long as the system is prepared in the state $|1; g\rangle$ initially. Due to the dissipation, the oscillation amplitude decreases as the time elapses and the system will eventually decay to the ground state $|0; g\rangle$. Furthermore, it is clear that, the life time of the cavity photon becomes much longer than the case with independent baths. From the above calculations, it is the coherence between the atom and the cavity field that slows down the decay of the system dramatically. The mechanism underlying the dramatic slowing decay will be explained later.

Vacuum Rabi splitting.— To further explore the effect of a common bath, we study the transmission spectrum of the system. To this aim, we drive the cavity by an external field. The action of the driven field on the cavity mode is described by

$$H_{driven} = \eta(a^\dagger e^{-i\omega_d t} + a e^{i\omega_d t}) \tag{7}$$

where η denotes the intensity of the driven field and ω_d its frequency.

The driven field induces the transition between the ground state and the excited states, while the dissipation causes the excited states to decay to the ground state. When the evolution time is long enough, the system will reach a steady state. To investigate the behavior of the steady state, we eliminate the time dependence from the Hamiltonian through the unitary transformation $U(t) = \exp[i(a^\dagger a + \sigma_z/2 + \sum_i b_i^\dagger b_i)\omega_d t]$. The atom-cavity Hamiltonian in the rotating frame becomes

$$\begin{aligned}\mathcal{H} &= i\frac{\partial}{\partial t}U(t)U^\dagger(t) + U(t)\left(H_{JC} + H_{driven}\right)U^\dagger(t) \\ &= \Delta_1 a^\dagger a + \frac{\Delta_2}{2}\sigma_z + g(a^\dagger\sigma^- + a\sigma^+) + \eta(a^\dagger + a)\end{aligned} \tag{8}$$

where $\Delta_{1(2)} = \omega_{c(0)} - \omega_p$ is the detuning between the cavity (atom) and the driven field. The last term can be regarded as a perturbation whenever $\eta \ll \omega_c, \omega_0, g$, i.e. weak driven field. Then the master equation can be written as

$$\frac{d}{dt}\rho_{cd} = -i\langle c[\mathcal{H}, \rho]|d\rangle + \sum_{k,l}\gamma^{cdkl}\rho_{kl}. \tag{9}$$

Therefore, the average photons number over the steady state is obtained numerically by finding the density matrix satisfying $d\rho_{cd}/dt = 0$ for any states $|c\rangle$ and $|d\rangle$. The results are shown in Fig. 3, where the average photon number reach its peaks when the driven field is just resonant with the energy difference between the dressed states $|1, \pm\rangle$ and the ground state $|G\rangle$. This effect is the vacuum Rabi splitting.

To our surprise, the double peaks are not symmetric if the atom-cavity coherence is taken into account. As shown in the figure, the peak corresponding to $\omega_d = E_1^- - E_0$ is elevated while the peak corresponding to $\omega_d = E_1^+ - E_0$ is suppressed compared with the case that neglects the effect of the coherence. Obviously this asymmetry is completely resulted from the coherence of the global system.

Physical mechanism.— Now let us explain the above two results in terms of the decay of the dressed states. Both of the atom and the cavity in their excited states will decay to their ground state due to the dissipation. Roughly speaking, the decay rate of the cavity mode is $J_1(\omega_c)$ and the spontaneous emission rate of the atom is $J_2(\omega_0)$. Since the two dressed states



FIG. 3. (Color online) The steady state of the system driven by an external field. The parameters are set as $\omega_0 = \omega_a = 1, g = 0.1, \omega_{c1} = 5, \omega_{c2} = 8, \alpha_1 = 0.002, \alpha_2 = 0.001, \eta = 0.005$.

$|1,\pm\rangle = (\pm|1;g\rangle + |0;e\rangle)/\sqrt{2}$ are symmetric and antisymmetric superpositions of excited atom and photon states, their decay rate would be $\Gamma_{1,\pm} = \left(\sqrt{J_1(\omega_c)} \pm \sqrt{J_2(\omega_0)}\right)^2$ instead of the sum of $J_1(\omega_c)$ and $J_2(\omega_0)$, the crossing terms proportional to $\sqrt{J_1(\omega_c)J_2(\omega_0)}$ comes from the effect of the atom-cavity coherence. Due to the coherence effect, the constructive interference between the atom channel and the cavity channel enhances the decay of the dressed state $|1,+\rangle$, while the destructive interference suppresses the decay of the dressed state $|1,-\rangle$.

	$\Gamma_{1\pm}$
secular appro	$\left(\sqrt{J_1(\omega_{1\pm,0})} \pm \sqrt{J_2(\omega_{1\pm,0})}\right)^2$
without coherence	$J_1(\omega_{1\pm,0}) + J_2(\omega_{1\pm,0})$
traditional	$J_1(\omega_c) + J_2(\omega_0)$

TABLE I. The decay rates of the dressed states obtained from different approaches.

A more accurate result for the decay rates of the dressed states $|1,\pm\rangle$ which are denoted by $\Gamma_{1,\pm}$ is shown in Table I, where we also give the results from that neglecting the coherence effect and the results from the master equation Eq. (2). As shown in the table, it satisfies $\Gamma_{1+}^{secular} > \Gamma_{1+}^{tra} = \Gamma_{1-}^{tra} > \Gamma_{1-}^{secular}$. The same conclusion can also be observed in Fig. 4, where we plot the probability for the system in the symmetric and antisymmetric dressed states, i.e. the time evolution of the diagonal elements ρ_{1+1+} and ρ_{1-1-}. It is clear that the symmetric state $|1,+\rangle$ decays more faster than the antisymmetry state $|1,-\rangle$. Apart from the numerical results, an analytical result based on iteration calculations and the comparison between the analytical and numerical approaches are shown in the supplemental material [27].

This result can be used to explain the slow decay of a cavity photon. Because the state of a single cavity photon can be regarded as a coherent superposition of the dressed state $|1,+\rangle$ and $|1,-\rangle$, the slow decay rate of a single cavity photon comes from that there is a component $|1,-\rangle$, which has a smaller decay rate.



FIG. 4. The probability for the system in its symmetric state $|1,+\rangle$ and antisymmetry state $|1,-\rangle$ in the absent of the external driven field. The parameters are set as $\omega_0 = \omega_a = 1$, $g = 0.1$, $\omega_{c1} = 5$, $\omega_{c2} = 8$, $\alpha_1 = 0.002, \alpha_2 = 0.001$.

Similarly, the asymmetric vacuum Rabi splitting can also be explained from the different decay rates of the states $|1,+\rangle$ and $|1,-\rangle$. Because the state $|1,-\rangle$ has a small decay rate, the steady state will have a larger component. This will give a stronger signal of the average photon number. In contrary, the state $|1,+\rangle$ has a larger decay rate, so the steady state will have a smaller component and it will give a weaker signal.

Conclusion.— We propose a scheme to realize the JCM with a common bath within the current experimental capabilities. Our results demonstrate how the coherence between two coupled subsystems affects the decay of the global system when they share a common bath by a simple JCM. The destructive interference between two decay channels weakens the decay of the antisymmetric state, while the constructive interference strengthes the decay of the symmetric state. Therefore, the antisymmetric state is more robust to the environment.

We hope the robustness of the antisymmetric state can be applied in quantum information process, such as the storage of the quantum state in quantum network. In addition, these common bath induced quantum coherence phenomena will promote the studies of fundamental problems of quantum open system.

Acknowledgments.— This work was supported by National NSF of China (Grant Nos. 10975181 and 11175247) and NKBRSF of China (Grant No. 2012CB922104).

---

* zhoudl72@aphy.iphy.ac.cn

[1] H. P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, Oxford, London, 2002).

[2] V. Braginsky and F. Khalili, *quantum measurement* (Cambridge University Press, London, 1992).

[3] G. M. Huang, T. J. Tarn, and J. W. Clark, J. Math. Phys. **24**, 2608 (1983).

[4] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, U.K., 2000).

[5] V. Ramakrishna and H. Rabitz, Phys. Rev. A **54**, 1715 (1996).

[6] A. J. Leggett, S. Chakravarty, A. T. Dorsey, M. P. A. Fisher, A. Garg, and W. Zwerger, Rev. Mod. Phys. **59**, 1 (1987).

[7] U. Weiss, *Quantum dissipative systems, third Edition*, (World Scientific Publishing, 2008).

[8] E. T. Jaynes and F. W. Cummings, Proc IEEE **51**,89 (1963).

[9] X. H. Bao, A. Reingruber, P. Dietrich, J. Rui, A. Duck, T. Strassel, L. Li, N. L. Liu, B. Zhao and J. W. Pan, Nature Phys. **8**, 512 (2012).

[10] T. Häyrynen, J. Oksanen, and J. Tulkki, Phys. Rev. A **83**, 013801 (2011).

[11] A. Stokes, A. Kurcz, T. P. Spiller, and A. Beige, Phys. Rev. A **85**, 053805 (2012).

[12] L. S. Bishop, E. Ginossar, and S. M. Girvin, Phys. Rev. Lett. **105**, 100505 (2010).

[13] M. Scala, B. Militello, A. Messina, J. Piilo, and S. Maniscalco, Phys. Rev. A **75**, 013811 (2007).

[14] M. J. Bhaseen, J. Mayoh, B. D. Simons, and J. Keeling, Phys. Rev. A **85**, 013817 (2012).

[15] B. Masashi, J. Phys. A: Math. Theor. **43**, 335305 (2010).

[16] G. Gangopadhyay, S. Basu, and D. S. Ray, Phys. Rev. A **47**, 1314 (1993).

[17] C. Cohen-Tannouji, J. Dupont-Roc, and G. Grynberg, *Atom-Photon Interactions: Basic Process and Applications*(John,Wiley & Sons,New York,1998)

[18] T. Hümmer, G. M. Reuther, P. Hänggi, and D. Zueco, Phys. Rev. A **85**, 052320 (2012).

[19] F. Nissen, S. Schmidt, M. Biondi, G. Blatter, H. E. Türeci, and J. Keeling, Phys. Rev. Lett. **108**, 233603 (2012).

[20] G. S. Agarwal, Phys. Rev. Lett. **57**, 1732 (1984).

[21] R. J. Thompson, G. Rempe, and H. J. Kimble, Phys. Rev. Lett. **68**, 1132 (1992).

[22] Y. F. Zhu, D. J. Gauthier, S. E. Morin, Q. L. Wu, H. J. Carmichael, and T. W. Mossberg, Phys. Rev. Lett. **64**, 2499 (1990).

[23] S. Savasta, R. Saija, A.Ridolfo, O. D. Stefano, P. Denti and F. Borghese, ACS. Nano. **4**, 6369 (2010).

[24] T. Yoshie, A. Scherer, J. Hendrickson, G. Khitrova, H. M. Gibbs, G. Rupper, C. Ell, O. B. Shchekin, and D. G. Deppe, Nature, **432**,200 (2004).

[25] H. Toida, T. Nakajima, and S. Komiyama, Phys. Rev. Lett. **110**, 066802 (2013).

[26] H. F. Wang, S. Ashhab, and F. Nori, Phys. Rev. A **83**, 062317 (2011).

[27] See Supplyment Materials for detailed derivation of the master equaiton and the analytical result for the matrix elements $\rho_{1+,1+}$ and $\rho_{1-,1-}$.